

N-dimensional alternate coined quantum walks from a dispersion relation perspective

Eugenio Roldán⁽¹⁾, Carlo Di Franco^(2,3), Fernando Silva⁽¹⁾, and Germán J. de Valcárcel⁽¹⁾,

⁽¹⁾*Departament d'Òptica, Universitat de València, Dr. Moliner 50, 46100-Burjassot, Spain*

⁽²⁾*Departamento de Química Física, Universidad del País Vasco-Euskal*

Herriko Unibertsitatea, Apartado 644, E-48080 Bilbao, Spain and

⁽³⁾*School of Mathematics and Physics, Queen's University, Belfast BT7 1NN, United Kingdom*

Abstract

We propose an alternative definition of an N-dimensional coined quantum walk by generalizing a recent proposal [Di Franco *et al.*, Phys. Rev. Lett. **106**, 080502 (2011)]. This N-dimensional alternate quantum walk, AQW_(N), in contrast with the standard definition of the N-dimensional quantum walk, QW_(N), requires only a coin-qubit. We discuss the quantum diffusion properties of AQW₍₂₎ and AQW₍₃₎ by analyzing their dispersion relations that reveal, in particular, the existence of diabolical points. This allows us to highlight interesting similarities with other well known physical phenomena. We also demonstrate that AQW₍₃₎ generates genuine multipartite entanglement. Finally we discuss the implementability of AQW_(N).

PACS numbers:

In both its standard forms, the coined [1] and the continuous one [2], the quantum walk is the quantum version of a classical random process, described by the diffusion and the telegrapher's equations, respectively [3]. In the coined quantum walk – the process we consider here – there is a system (the walker) that undergoes a conditional displacement, to the right or left, depending on the output of a *coin throw*, as in the random walk. But differently from its classical counterpart, here both coin and walker are quantum in nature; hence the walker position coherently spans over the lattice sites. The one-dimensional coined quantum walk – QW₍₁₎ for short – has been studied from many different perspectives, especially from the quantum computational point of view [4]. In the last few years it has also received increasing experimental attention [5–7], including the case of quantum walks with more than one particle [8].

The situation is quite different when dealing with N-dimensional quantum walks, QW_(N) for short. They were first discussed by Mackay *et al.*, who introduced them in complete analogy with QW₍₁₎ [9] (see also Ref. [10]). As defined in Ref. [9], QW_(N) requires the use of a 2N-dimensional qudit as coin, as well as a coin operator represented by a $2N \times 2N$ unitary matrix. This introduces increasing complexity in the process as N grows, especially from the experimental viewpoint [11, 12], but also from the theoretical one [13, 14]. However, Di Franco *et al.* [15] have recently proposed an alternative two-dimensional quantum walk, namely the *alternate* quantum walk – AQW for short – that is simpler than the standard one. In AQW the coin is a single qubit, as in QW₍₁₎, and each time step is divided into two halves: in the first one the coin is thrown (i.e., a Hadamard transformation is applied on the coin-qubit) and the conditional displacement along \vec{x} is consequently performed; then, in the second half of the time step, the coin is thrown again and the conditional displacement along \vec{y} is performed. Hence, in AQW, the four-dimensional qudit of QW₍₂₎ is

replaced by a single qubit, the price paid for that being to double the number of sub-steps per single time step. Quite unexpectedly, AQW reproduces the same spatial probability distributions of QW₍₂₎ when the Grover coin is used – Grover-QW₍₂₎ for short – for a set of particular initial conditions, precisely those for which the characteristic localization of Grover-QW₍₂₎ does not occur. In Refs. [15, 16] analytical demonstrations of the (limited) equivalence between Grover-QW₍₂₎ and AQW are given.

Here we generalize AQW to N-dimensions: We define the alternate N-dimensional coined quantum walk – AQW_(N) for short – as a quantum walk in which the time steps are divided into N sub-steps. In each of these sub-steps, the coin throw is followed by the conditional displacement along one of the N dimensions. In a single time step of AQW_(N), the qubit-coin is therefore thrown N times, but this is clearly simpler than throwing a single 2N-dimensional coin, from both the experimental and analytical points of view, in particular for large N. We show below that, besides a simpler experimental implementability as compared with QW_(N), AQW_(N) has a rich dynamics that is also simpler to understand than that of QW_(N). We provide some analytical results concerning the evolution of the probability distribution for cases $N = 2, 3$, paying special attention to the processes' dispersion relations (that reveal the existence of diabolical points), and demonstrate the existence of genuine multipartite entanglement in AQW₍₃₎. We end this letter with a brief discussion concerning the experimental implementability of AQW_(N).

In order to introduce AQW_(N) formally, let $|\psi\rangle_t$ represent the state of the system at (discrete) time t . Vector $|\psi\rangle_t$ is defined in the compound Hilbert space $\mathcal{H}_C \otimes \mathcal{H}_P$, where \mathcal{H}_C and \mathcal{H}_P are the Hilbert spaces for the coin-qubit and lattice sites, respectively. Being \mathcal{H}_C and \mathcal{H}_P spanned, respectively, by $\{|c\rangle, c = u, d\}$ and

$\{|\vec{x}\rangle, \vec{x} \in \mathbb{Z}^N\}$, we can write

$$|\psi\rangle_t = \sum_{c=u,d} \sum_{\vec{x}} c_{\vec{x},t} |\vec{x}; c\rangle, \quad (1)$$

$|\vec{x}; c\rangle = |\vec{x}\rangle \otimes |c\rangle$, where $c_{\vec{x},t}$ is the probability amplitude for the walker to be at site $\vec{x} = (x_1, \dots, x_N)$ at time t with the coin in state c . The probability of finding the walker at site \vec{x} and time t is $P_{\vec{x},t} = |u_{\vec{x},t}|^2 + |d_{\vec{x},t}|^2$.

The state evolves as $|\psi\rangle_{t+1} = \hat{U}_{(N)} |\psi\rangle_t$, with $\hat{U}_{(N)} = \hat{D}_N \hat{C}_N \hat{D}_{N-1} \hat{C}_{N-1} \dots \hat{D}_1 \hat{C}_1$ a unitary operator. Here, operator \hat{C}_i is the *coin operator*, acting only in \mathcal{H}_C , whose more general form is

$$\begin{aligned} \hat{C}_i = & \cos \theta_i \left(|u\rangle \langle u| - e^{i(\alpha_i + \beta_i)} |d\rangle \langle d| \right) \\ & + \sin \theta_i \left(e^{i\alpha_i} |u\rangle \langle d| - e^{i\beta_i} |d\rangle \langle u| \right), \end{aligned} \quad (2)$$

with $(\alpha_i, \beta_i, \theta_i)$ arbitrary reals and $i = 1, \dots, N$ (notice that, for $\alpha_i = \beta_i = 0$ and $\theta_i = \pi/4$, \hat{C}_i is just the Hadamard transformation). Operator \hat{D}_i is the conditional displacement operator along direction x_i , which we write as

$$\hat{D}_i = \sum_{\vec{x} \in \mathbb{Z}^N} [|\vec{x} + \vec{n}_i; u\rangle \langle \vec{x}; u| + |\vec{x} - \vec{n}_i; d\rangle \langle \vec{x}; d|], \quad (3)$$

where \vec{n}_i is the unit vector along direction x_i .

Equation $|\psi\rangle_{t+1} = \hat{U}_{(N)} |\psi\rangle_t$ can be expressed as a two-dimensional map relating probability amplitudes $c_{\vec{x},t+1}$ with $\tilde{c}_{\vec{x}',t}$ with $c, \tilde{c} = u, d$ and \vec{x}' nearest neighbors to \vec{x} . This map admits two plane-wave solutions of the form $\text{col}(u_{\vec{q},t}, d_{\vec{q},t})_{\pm} = \vec{\varphi}_{\vec{q},\pm} \exp[i(\vec{q} \cdot \vec{x} - \omega^{(\pm)} t)]$ where $\vec{\varphi}_{\vec{q},\pm} = \text{col}(u_{\vec{q},\pm}, d_{\vec{q},\pm})$ are time independent vectors, \vec{q} is the pseudo-momentum, with $q_i \in (-\pi, \pi]$, and $\omega^{(\pm)}$ are two frequencies determined by the dispersion relation.

The dispersion relation is most relevant because $|\psi\rangle_t$ is entirely determined by it, given $|\psi\rangle_{t=0}$. Moreover, when the initial state extends over a finite set of points in the lattice, especially when it is modulated by a *smooth* function of space, the dispersion relation is particularly useful for predicting the evolution of the initial wave-packet, due to the relatively well defined group velocity (given by the local gradient of the dispersion relation curve). In this case, long-wavelength continuous approximations are very well suited and useful for envisaging the long time behavior of the probability distribution. This has been discussed in detail for QW₍₁₎ in Ref. [18] (see also Refs. [3, 17]). Here, we will limit ourselves to a qualitative discussion of what the dispersion relation suggests for cases $N = 2, 3$.

Let us first consider AQW₍₂₎. By proceeding as stated above, one obtains the following dispersion relation

$$\cos \Omega = c_1 c_2 \cos(u + v) + s_1 s_2 \cos(u - v), \quad (4)$$

where $c_i = \cos \theta_i$, $s_i = \sin \theta_i$, $u = q_1 + (\beta_1 + \alpha_2)/2$, $v = q_2 + \beta_2/2$, and $\Omega = \omega - (\beta_1 + \beta_2 + \alpha_2)/2$. Notice

that phase α_1 does not appear in Eq. (4), hence it is irrelevant. As for the phases that do appear, they just entail a translation of the frequency and spatial quasi-momentum. Only θ_1 and θ_2 are dynamically relevant parameters.

In Fig. 1(a) we have represented the dispersion curves for $\theta_1 = \theta_2 = \pi/4$. The most relevant features are the existence of a number of saddle-points (for which the group velocity is zero), together with regions of maximum slope, which equals 0.5 (hence the maximum velocity in AQW₍₂₎) and, most importantly, the existence of *diabolical points* (DPs in the following). DPs are conical intersections involving a degeneracy [20]. Interestingly, when $\theta_1 \neq \theta_2$, the DPs disappear, as shown in Fig. 1(b). This suggests that the existence of DPs could be particularly sensitive to decoherence effects in the coin mechanism.

The dynamics around the points of null and maximum slope can be analyzed as in Ref. [18] for QW₍₁₎, i.e., in terms of known solutions of simple linear wave equations. One can then envisage more or less straightforward generalizations of the results there discussed to the two-dimensional case (we discuss this elsewhere for Grover-QW₍₂₎ [22]). But the existence of DPs is particularly appealing and constitutes a qualitative difference with the one-dimensional case. This geometric object, the DP (that takes its name from the diabolo-like shape of the conical intersection), appears in physics in quite different contexts like, for instance, the electronic spectrum of polyatomic molecules [19], quantum triangular billiards [20], conical refraction in crystal optics [21], or the dispersion relations for massless fermions (Dirac electrons) in QED and for electrons in graphene [23, 24], and the diabolo is associated with some of the remarkable phenomena appearing in those systems. We can expect to find in the dynamics of AQW₍₂₎ some parallelism with phenomena present in the aforementioned systems. In this sense, Fig. 1(c) shows the propagation of an initially extended state (with a Gaussian probability distribution of $\sigma_{HWHM} = 7$) after 90 time steps for an initial coin state $\text{col}(1, i)/\sqrt{2}$ equal for all the populated sites. A homogeneous ballistic propagation from the origin with perfect circular symmetry is clearly visible, which recalls the ring shape of the beam exiting a biaxial crystal under conical refraction conditions [21]. It is interesting to compare this dynamics with that shown in Fig. 1(d), which has been obtained for the same initial conditions but with $\theta_1 = \pi/4 \neq \theta_2 = \pi/3$. In this case the diabolo is lost and the branches of the dispersion relation show a parabolic shape [see Fig. 1(b)] that leads to evolutions typical of linear optical diffraction [18]. It is interesting to mention that a similar controlled disappearance of the DP has been experimentally observed in graphene [24].

Before moving to a higher dimension, let us revisit the relation between AQW₍₂₎ and Grover-QW₍₂₎ from the dispersion relation perspective. The dispersion relations for Grover-QW₍₂₎ consist of four sheets, because

the coin space is four-dimensional, and can be found in Refs. [14, 22]. In our notation, they read $\omega_{1,2} = 0, \pi$ and $\omega_{3,4} = \pm \arccos[(\cos u + \cos v)/2]$. Remarkably, the two sheets $\omega_{3,4}$ coincide with those of $\text{AQW}_{(2)}$, Eq. (4), for $\theta_1 = \theta_2 = \pi/4$ after identifying (u, v) in $\text{Grover-QW}_{(2)}$ with $(u + v, u - v)$ in $\text{AQW}_{(2)}$, i.e., both dispersion relations coincide for these parameters up to a $\pi/4$ rotation of the pseudo-momentum. The two other roots in $\text{Grover-QW}_{(2)}$, $\omega_{1,2} = 0, \pi$, are constant, which means that the projections of the initial state onto the corresponding eigenvectors will not evolve in time. This is the origin of localization in $\text{Grover-QW}_{(2)}$ for most initial coin states, as already noticed in Ref. [14]. We conclude that, whenever the initial state in $\text{Grover-QW}_{(2)}$ does not project onto the eigenvectors governed by $\omega_{1,2}$, $\text{Grover-QW}_{(2)}$ and $\text{AQW}_{(2)}$ are isomorphous for $\theta_1 = \theta_2 = \pi/4$. This is our proof of the (partial) equivalence between the two versions of the process.

Let us now move to $\text{AQW}_{(3)}$. The dispersion relation is governed by

$$\begin{aligned} \sin \Omega = & c_1 [c_2 c_3 \sin(u + v + w) + s_2 s_3 \sin(u - v + w)] \\ & + s_1 [c_2 s_3 \sin(u + v - w) - s_2 c_3 \sin(u - v - w)], \end{aligned} \quad (5)$$

where $c_i = \cos \theta_i$, $s_i = \sin \theta_i$, and $(u, v, w, \Omega) = (q_1, q_2, q_3, \omega) + (\delta q_1, \delta q_2, \delta q_3, \delta \omega)$. Here $2\delta q_1 = \alpha_1 + \beta_2$, $2\delta q_2 = \alpha_2 + \beta_3$, $2\delta q_3 = \alpha_3 + \beta_1$, and $\delta \omega = -(\delta q_1 + \delta q_2 + \delta q_3)$. From (5) two dispersion relations are obtained, namely $\omega^{(+)} = \Omega$ and $\omega^{(-)} = \pi - \omega^{(+)}$. As in $\text{AQW}_{(2)}$, some phases in \hat{C}_i are absent in the dispersion relation (hence they are irrelevant) and the effect of the rest of phases in \hat{C}_i is just a displacement of the dispersion relation surfaces in the $\{q_1, q_2, q_3, \omega\}$ space. Eq. (5) is simpler for $\theta_i = \pi/4$, $i = 1, 2, 3$

$$\sin \omega = \frac{1}{\sqrt{2}} [\sin v \cos(u - w) + \cos v \sin(u + w)], \quad (6)$$

in which case there are eight degeneracies (occurring when $\omega^{(+)} = \omega^{(-)} = \pi/2$) at $(u_{DP}, v_{DP}, w_{DP}) = \{(a, a, a), (-a, -a, b), (-a, b, -a), (b, -a, -a)\}$, with $a, b = \pi/4, 3\pi/4$ and $a \neq b$. These are the 3D equivalents of the DPs discussed for $\text{AQW}_{(2)}$. As in $\text{AQW}_{(2)}$, the DPs disappear when phases θ_i are different. We mention that several crystallographic structures have recently been proposed for obtaining 3D-DPs (Dirac-semimetal in 3D) [25].

In Figs. 2(a,b) we represent two bidimensional projections of the probability distribution corresponding to the propagation of a walker initially localized at the origin. The width of the distribution grows linearly with time, along all the three spatial dimensions, as it happens for lower dimensionality. In order to show the effect of the DP, Figs. 2(c,d) show the same projections as Figs. 2(a,b) when the initial condition is not localized

but extended. Again we choose a Gaussian distribution with $\sigma_{HWHM} = 7$, whose pseudo-momentum is centered at one of the DPs. We observe a symmetric ballistic dynamics in the (x_1, x_2) plane, Fig. 2(c), that resembles that of Fig. 1(c). Notice the existence of two concentric bright rings, as it occurs in conical refraction [20]. However, the probability is not equally symmetric in the (x_1, x_3) plane, Fig. 2(d), which reveals an intrinsic lack of symmetry in $\text{AQW}_{(3)}$: indeed it can be shown that it is not possible to find any initial coin state that leads to a symmetric propagation in all directions (the chosen initial coin state, $(0, 1)$ for all the initially populated sites, leads to a symmetric distribution in the planes with constant x_3 , but not in those for which x_3 varies, as the figures show).

One could wonder whether $\text{AQW}_{(3)}$ is a process similar to $\text{Grover-QW}_{(3)}$ as it happens for $N = 2$. The answer is negative: we have derived and compared the dispersion relations of both processes for $N = 3$ and they are different. Hence $N = 2$ is a singular case in this respect.

Having a richer and more complex structure than the bipartite case [26], multipartite entanglement has recently attracted a lot of interest in the scientific community. Clearly, a feasible system able to generate a proper amount of genuine multipartite entanglement could be a valuable benchmark for this rapidly growing research field. Hence a most relevant question is whether quantum walks with $N > 2$ do exhibit genuine multipartite entanglement or not. We have investigated this for $N = 3$. For analyzing its generation in $\text{AQW}_{(3)}$, we have first to trace out the state of the coin. We are then left with a density matrix in the composite Hilbert space $\mathcal{H}_{x_1} \otimes \mathcal{H}_{x_2} \otimes \mathcal{H}_{x_3}$ (each subspace corresponding to a direction of the walk). We evaluate the multipartite entanglement present in this composite system by means of the tripartite negativity [27]. This is defined as the geometric average of the three negativities that are obtained by considering the three possible bipartitions of the total system, giving

$$N^{(3)} = \sqrt[3]{N_{1-23} N_{2-13} N_{3-12}}. \quad (7)$$

Here N_{i-jk} is the negativity of the composite system $\{i, j, k\}$ corresponding to the bipartition in the subsystem $\{i\}$ and the subsystem $\{j, k\}$. Each Hilbert direction-subspace has a dimension growing with the number of time steps, so we use the generalization of the negativity for higher-dimensional systems (so as to have $0 \leq N \leq 1$) [28]. We have evaluated $N^{(3)}$ in $\text{AQW}_{(3)}$, with the walker starting at the origin and initial coin state $\text{col}(1, i)/\sqrt{2}$, for a number of time steps t up to 10, obtaining the plot in Fig. 2(e). Even if the number of time steps considered here is not so large (due to the dimension of the total Hilbert space, the computational power required for the evaluation grows rapidly with t), it is easy to check that $\text{AQW}_{(3)}$ is able to properly generate genuine multipartite entanglement.

Let us finally discuss the implementability of $AQW_{(N)}$. Realizing $QW_{(N)}$ is quite demanding because of the complexity of implementing coin-operators to transform the needed $2N$ -dimensional qudit. On the other hand, in $AQW_{(N)}$: (i), a single coin-qubit is required independently of N ; (ii), two-dimensional transformations of the qubit are easy to implement [5–7]; and (iii), the sequential application of operators $\hat{D}_j\hat{C}_j$, $j = 1, \dots, N$, makes the implementation of $AQW_{(N)}$ similar to that of $QW_{(1)}$ provided that all N dimensions could be multiplexed into a single one (similarly to what Schreiber *et al.* [12] have recently done in their pioneering implementation of $QW_{(2)}$). Indeed one could even implement $AQW_{(N)}$ with a constant number of physical elements independently of N if one has sufficient control on the experimental device.

In order to illustrate the idea, we generalize the idealized device already discussed in Refs. [17, 29], which is similar to that actually used in Ref. [5]. Consider a long enough optical cavity containing two electro-optic modulators (EOM) whose roles are (i), EOM1 performs the coin-operator \hat{C} (i.e., makes a unitary transformation of the light polarization state, which plays the role of the coin-qubit in this implementation of the walk); and (ii), EOM2 performs the conditional displacement \hat{D} , which consists in up/down shifting the carrier frequency of the light pulse depending on its polarization. The light pulses entering the cavity are assumed to be much shorter than the cavity length, and the frequency shifts introduced by EOM2 must be large enough for avoiding any frequency overlapping between pulses. With such a device not only $QW_{(1)}$ can be implemented, see [5, 17, 29], also $AQW_{(N)}$ could be implemented for different values of N by properly programming the operation of the EOMs, without the need for additional elements. For example, in order to perform $AQW_{(2)}$, the first half of the time step is implemented during one cavity round-trip of the light pulse (during which EOM1 and EOM2 implement $\hat{D}_1\hat{C}_1$); then, in the subsequent cavity round-trip, the settings of both EOMs are changed in order to perform a different coin operator and a different frequency displacement implementing $\hat{D}_2\hat{C}_2$. Importantly, the frequency displacements in \hat{C}_1 and \hat{C}_2 must be different enough in order to multiplex a large number of steps [12]. Only technical limitations seem to limit the extension of the procedure to higher N . In any case, with this oversimplified discussion, we are not claiming that the device just outlined is the most appropriate for implementing $AQW_{(N)}$. Indeed a suitable modification of the flexible device used by Schreiber *et al.* [7, 12] would probably be a more promising option. With our discussion we just want to emphasize that a single and conceptually simple device *could* implement alternate quantum walks with tunable dimensionality. It is much interesting that such a device would be a source of genuine multipartite entanglement.

In conclusion, we have introduced the N -dimensional alternate quantum walk and discussed some of its proper-

ties through the analysis of the dispersion relations that reveal, in particular, the existence of diabolical points. We have also demonstrated that for $N = 3$ the process generates genuine multipartite entanglement. We have finally discussed the implementability of the process that could be possible with physical resources that do not necessarily grow with the dimensionality of the walk.

We thank Th. Busch, M. Mc Gettrick, C. M. Chandrashekar, C. Navarrete-Benlloch, A. Pérez, A. Romanelli, and M. Hinarejos for discussions. This work has been supported by the Spanish Government and the European Union FEDER through Projects FIS2008-06024-C03-01 and FIS2011-26960, and by the Basque Government through Grant IT472-10. C.D.F. acknowledges A. Pérez and the Universitat de València for the kind hospitality.

-
- [1] Y. Aharonov, L. Davidovich, and N. Zagury, Phys. Rev. A **48**, 1687 (1993); D. Meyer, J. Stat. Phys. **85**, 551 (1996); J. Watrous, Proc. 33rd Symposium on the Theory of Computing (ACM Press, New York, 2001), p.60.
 - [2] E. Farhi and S. Gutman, Phys. Rev. A **58**, 915 (1998); A.M. Childs and J. Goldstone, Phys. Rev. A **70**, 042312 (2004).
 - [3] F.W. Strauch, Phys. Rev. A **73**, 054302 (2006).
 - [4] A. Ambainis, Int. J. Quant. Inf. **1**, 507 (2003); J. Kempe, Contemp. Phys. **44**, 307 (2003); V. Kendon, Math. Struct. Comp. Sci. **17**, 1169 (2006); V. Kendon, Phil. Trans. R. Soc. A **364**, 3407 (2006); N. Konno, *Quantum Walks*, in Quantum Potential Theory, Lect. Notes Math., Vol. 1954, edited by U. Franz and M. Schürmann (Springer, 2008).
 - [5] D. Bouwmeester et al., Phys. Rev. A **61**, 013410 (2000)
 - [6] B. Do et al. J. Opt. Soc. Am. B **22**, 499 (2005); C. A. Ryan et al., Phys. Rev. A **72**, 062317 (2005); P. Zhang et al., *ibid* **75**, 052310 (2007); P. H. Souto Ribeiro et al., *ibid* **78**, 012326 (2008); M. Karski et al., Science **325**, 174 (2009); H. B. Perets et al., Phys. Rev. Lett. **100**, 170506 (2008); H. Schmitz et al., *ibid* **103**, 090504 (2009); F. Zähringer et al., *ibid* **104**, 100503 (2010).
 - [7] A. Schreiber et al., Phys. Rev. Lett. **104**, 050502 (2010); *ibid* **106**, 180403 (2011).
 - [8] A. Peruzzo, et al., Science **329**, 1500 (2010); J. C. F. Matthews, et al., arxiv: quant-ph/1106.1166 (2011); L. Sansoni, et al., Phys. Rev. Lett. **108**, 010502 (2012).
 - [9] T.D. Mackay, S.D. Bartlett, L.T. Stephenson, and B.C. Sanders, J. Phys. A: Math. Gen. **35**, 2745 (2002).
 - [10] C. Moore and A. Russell, *Proceedings of RANDOM, 2002*, edited by J.D.P. Rolim and P. Vadham (Springer, Cambridge, MA, 2002), pp.164-178.
 - [11] E. Roldán and J.C. Soriano, J. Mod. Opt. **52**, 2649 (2005).
 - [12] A. Schreiber et al. arxiv: quant-ph/1204.3555 (2012).
 - [13] B. Tregenna et al., New. J. Phys. **5**, **83** (2003); I. Carneiro et al., New J. Phys. **7**, 156 (2005); A.C. Oliveira et al., Phys. Rev. **74**, 012312 (2006); K. Watabe et al., Phys. Rev. A **77**, 062331 (2008); M. Stefanák et al., Phys. Scr. **T140**, 014035 (2010); N. Shenvi et al., Phys. Rev. A **67**,

- 052307 (2003).
- [14] N. Inui, Y. Konichi, and N. Konno, Phys. Rev. A **69**, 052323 (2004).
- [15] C. Di Franco, M. Mc Gettrick, and Th. Busch, Phys. Rev. Lett. **106**, 080502 (2011).
- [16] C. Di Franco, M. Mc Gettrick, T. Machida, and Th. Busch, Phys. Rev. A **84**, 042337 (2011).
- [17] P.L. Knight, J.E. Sipe, and E. Roldán, Phys. Rev. A **68**, 020301 (2003); J. Mod. Opt. **51**, 1761 (2004).
- [18] G.J. de Valcárcel, E. Roldán, and A. Romanelli, New J. Phys. **12**, 123022 (2010).
- [19] C. A. Mead and D.G. Truhlar, J. Chem. Phys. **70**, 2284 (1979); L.S. Cederbaum et al., Phys. Rev. Lett **90**, 013001 (2003).
- [20] M. V. Berry and M. Wilkinson, Proc. Roy. Soc. A **392**, 15 (1984).
- [21] M.V. Berry and M.R. Jeffrey, Prog. Opt. **50**, 13 (2007).
- [22] M. Hinarejos, A. Romanelli, A. Pérez, E. Roldán, and G.J. de Valcárcel, to be published.
- [23] A.H. Castro Neto et al., Rev. Mod. Phys. **81**, 109 (2009).
- [24] A. Bostwick et al., New J. Phys. **9**, 385 (2007).
- [25] S.M. Young et al. Phys. Rev. Lett. **108**, 140405 (2012).
- [26] R. Horodecki et al., Rev. Mod. Phys. **81**, 865 (2009).
- [27] C. Sabin and G. Garcia-Alcaine, Eur. Phys. J. D **48**, 435 (2008).
- [28] S. Lee et al., Phys. Rev. A **68**, 062304 (2003).
- [29] P.L. Knight, J.E. Sipe, and E. Roldán, Opt. Commun. **227**, 147 (2003).
- [30] The multiplexing requires that the frequency displacements corresponding to different dimensions be very different (in fact this difference determines the maximum number of steps that can be implemented, see the discussion in [12]). Moreover, the frequency shifts should be done with a high repetition rate (larger than the inverse of the cavity round-trip time). All this could be very demanding for actual EOMs.

Figure captions

Fig. 1. 3D views of the two branches of (4), for $\theta_1 = \theta_2 = \pi/4$ (a) and $\theta_1 = \pi/4$ and $\theta_2 = \pi/3$ (b). Propagation in AQW₍₂₎ after 90 time steps of an initial state with a Gaussian probability distribution of width $\sigma_{HWHM} = 7$ and initial coin state $\text{col}(1/\sqrt{2}, i/\sqrt{2})$ equal for all populated sites, for $\theta_1 = \theta_2 = \pi/4$ (c) and $\theta_1 = \pi/4$ and $\theta_2 = \pi/3$ (d).

Fig. 2. Propagation in AQW₍₃₎ after 90 time steps of (a,b) a localized initial state, and (c,d) a spatially extended initial state (Gaussian probability distribution of $\sigma_{HWHM} = 7$) for $\theta_1 = \theta_2 = \pi/4$ and initial coin state $\text{col}(0, 1)$. In (e) the tripartite negativity is shown as a function of the number of steps.